

Valorisation of microbial biodiversity: Characterization of a milk-clotting enzyme produced by *Bacillus velezensis* FK6A strain isolated from Algerian dairy farm soil in light of its use in the manufacture of hard cooked cheeses.

F. Boumediene¹, S. Talantikite², S. Haddad³, A. Nouani^{2*}, S. Triki³, S. Belbraouet

¹University of Yahia Fares (Médeä)

²Laboratory of Food Technology and Human Nutrition, High National School of Agronomy, BP 16002, El Harrach, Algiers, ALGERIA University of Boumerdes, Food

³Technology Laboratory, BP 35000, Boumerdes, ALGERIA

*Corresponding author: a.nouani@univ-boumerdes.dz – Tel:(213)552579680

ARTICLE INFO

Article History :

Received :02/02/2020
Accepted :26/07/2020

Key Words:

Bacillus velezensis;
characterization;
culture medium;
identification;
Milk clotting enzyme;

ABSTRACT/RESUME

Abstract: A strain known as FK6A, which produces a milk-clotting enzyme isolated from the soil of an Algerian dairy factory, was identified as *Bacillus velezensis* based on morphology and internal transcription spacer sequence. Phylogenetic and sequencing analysis of 16S rDNA gene showed a 100% identity match between the tested strain and the *Bacillus velezensis* in the database (Code MH368142). This strain produced an active extracellular protease purified by partial purification using ammonium sulfate fractionation (40–80% saturation) followed by chromatography of molecular exclusion, which revealed the existence of two active peaks. The optimal activity was observed at pH 5.7; 75°C and 40 mM of CaCl₂. The clotting enzyme has been shown to be a metalloprotease inhibited by EDTA. The highly proteolytic activity of the *Bacillus velezensis* extracts observed on the curds appears a few minutes after the appearance of a firm coagulum reflected by the appearance of an abundant serum. Thus, to solve this problem and to inhibit proteolytic activity, the curds were cooked after brewing during the manufacturing process.

I. Introduction

Obtaining natural milk coagulants is a challenge, because the supply of rennet is limited compared to the dairy industry's rising demand for this enzyme. Natural coagulants of animal origin are costly, and for religious or dietary purposes, their use has been limited. This scenario has led to increased demand for new rennet replacements, promoting a search for new sources of proteases with coagulant properties [1, 2].

Microbial milk-clotting enzymes (MCE) have certain benefits over those derived from plants or animals, such as lower costs, greater biochemical

diversity and easier genetic modification. Some MCE development work has now centered on solid state fungal fermentation such as *Mucor miehei*, *Rhizomucor pusillus* var., *Aspergillus oryzae* are already widely used in commercial cheese production [1]. Otherwise, as possible rennet alternatives, certain bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, and *Enterococcus faecalis* have been suggested [3]. Singh et al [4] and McGugan [5] suggested that bacterial MCE have contributed to distinguishing flavours in Cheddar cheese

production. In addition, different sources of microorganisms make a variety of cheeses with and organoleptic properties [6, 7]. The aim of this study was to elucidate the nature and characteristics of the novel milk clotting enzyme produced by FK6A (a soil borne *Bacillus velezensis* strain). The conditions for the fermentation of the solid state and the synthesis of enzymes were investigated and trials for the production of baked cheese are carried out.

II. Material and Methods

II. 1. Microorganism, culture conditions and isolation method

Fifty grammes of soil samples were collected near a dairy factory in the Boumerdes region (about 50 km to the north of Algeria). Under magnetic agitation, a small amount of soil was homogenized with 90 mL sterile physiological water for 20 min to create an initial dilution (10^{-1}). Sample dilutions were made for each sample, and 0.1 mL of the correct 10^{-9} dilution was applied to the plate count agar (PCA) medium containing 2% (w/v) actidione antifungal agent to prevent fungal growth and 12% (w/v) sterilized skimmed milk powder. For the enumeration of the caseolytic strain developing milk clotting protease the plates were incubated at 30°C for 48 h by the presence of a strong halo around the colony. Colonies with distinct differences in morphology such as color (Gram technique), shape, and size were selected and purified using the same medium to streak. All isolates were tested for the development of milk clotting, and microscopically examined before stock preparations. The gram-positive bacterial isolates known as *Bacillus* have been purified, and the selected strains displaying coagulation activities have been preserved on agar conservation.

II. 2. Identification of FK6A strain

Talantikite et al [8], recorded the same strain identification method. Under optimized conditions, the PCR products were sequenced by Geno screen (Geno-screen Campus, Institut Pasteur de Lille 1, France) using Sanger technique. Since the sequencing results were collected, the 16S rDNA nucleotide sequence (724 bp sequence) of the strain was deposited in GenBank and matched with the 16S rDNA sequences available in the National Center for Biotechnology Information (NCBI) public databases (<http://www.ncbi.nlm.nih.gov/>) and LeBibi database (<http://umr5558-sud-str1.univ-lyon1.fr/lebibi/lebibi.cgi>) that uses the simple local alignment search software (BLAST).

II. 3. Protease production in shake flask cultures medium

different rheological

II. 3.1. Protease production in shake flask cultures, medium composition and inoculum:

For fermentation, basal medium (pH 6.8) used in the milk clotting protease production, containing (g/L): wheat bran (50), yeast extract (3), glucose (4), casein (2), and Na_2HPO_4 (3). The medium (50 mL) placed in an Erlenmeyer 250 mL flask was inoculated with 15×10^8 CFU/mL (Mac Farland 4.42) and incubated for 24 h in a control shaker at 30°C on a rotary shaker with agitation (110 rpm). The contents of the incubated Erlenmeyer flasks are first washed onto a gaze band under a sterile host, placed on a funnel, to recover the crude enzyme extract. At 4°C, the filtrate then undergoes centrifugation at 4000 rpm for 30 min. [9] The supernatant recovered forms the crude enzymatic extract.

II. 4. Analytical procedure

II. 4. 1. Milk clotting activity

Milk clotting activity (MCA) was measured with slight modifications of the procedure described by Shieh et al [10]. Skim milk powder was reconstituted by dissolving 12.5 g CaCl_2 solution in 100 mL of 0.5 g/L (pH 6.5). The clotting assay was performed by mixing 10 mL of this substrate and 1 mL of sample which is based on the visual assessment of the first clotting flakes' appearance. The clotting activity was expressed as Soxhlet Units (SU). One SU is defined as the amount of enzyme that coagulates 1 mL of a solution that contains 0.12 g skim milk powder and 0.11 mg CaCl_2 at 35°C in 40 min. The clotting activity was determined with the following formula:

$$US = 2400 \times 10 \times D/T$$

Where: T =clotting time and D =dilution of crude extract.

II. 4. 2. Proteolytic activity assay

Proteolytic activity was determined using the method reported by Silva et al [11] and Shieh et al [10]. The reaction mixture consisted of 0.4 mL 0.5% casein (Sigma) (w/v) in distilled water, 0.4 mL of 0.2 mol/L acetate buffer at pH 5.5, and 0.2 mL of crude enzyme. The reaction occurred in a bath at 35°C and at 30 min, the reaction was stopped by adding 1 mL of 10% trichloroacetic acid (TCA). 2 mL of the above filtrate was added by 5 mL NaOH (0.28 N) solution and 1.5 mL Folin-Ciocalteu phenol solution: water (1:2). After the mixture was kept at 35°C for 15 min, optical density (OD) at 660 nm was measured. The proteolytic activity was expressed in terms of units corresponding to $\text{OD}_{660\text{nm}}$ (UV-Visible Spectrophotometer, model V560, Shimadzu, Japan).

The ratio (R) was determined by dividing the milk clotting activity (MCA) by the proteolytic activity

II. 4.3. Protein concentration assay

The protein content was determined according to Lowry method [12] using crystalline bovine albumin as standard.

II. 5. Purification of the crude enzymatic extract

II. 5. 1. Partial purification of the milk-clotting enzyme (NH₄)₂SO₄ fractionation

The crude enzyme extract is exposed to an ammonium sulphate precipitation at three stage saturation (w/v): 40%, 60%, 80%. According to the 4°C ammonium sulphate precipitation table, 4.86 g, 7.80 g and 11.22 g are added to 20 mL of enzymatic extract. The precipitate collected at 10,000 g at 4°C for 20 min after centrifugation was suspended in the acetate buffer (0.01 M, pH5.5) and dialyzed overnight against repeated adjustments of the same buffering.

II. 5. 2. Size-exclusion chromatography

The concentrated extract was pre-equilibrated with an acetate buffer (0.01 M, pH 5.5) on the Sephadex G-75 column (30 cm x 1.5 cm). The 1.5 mL enzyme fractions were eluted with the same buffer at 36 mL/h and tested for enzyme activity and protein content. To further analysis, active fractions of the enzymes were pooled and lyophilized.

II. 5. 3. Molecular weight determination

The molecular weight of the protease from *B. velezensis* FK6A was determined upon purification of the active fraction on Sephadex G-75. The column was measured using a mixture of established molecular weight

standard proteins

including BSA (67 kDa), ovalbumin (40kDa), carbonic anhydrase (30kDa), and trypsin (23.8kDa).

II. 5. 4. Effect of inhibitors on enzyme activity

The effect of the enzyme inhibitors was evaluated using the Raposo and Domingos [13] technique. The residual activity was calculated after preincubation of the crude enzyme extract at 37°C for 30 min with inhibitors under the conditions defined by He et al [14]. Different protease inhibitors including

by the formula $R = \text{MCA (SU)}/\text{PA (POD}_{660\text{nm}})$.

particularly a metalloprotease inhibitor (ethylenediaminetetraacetic acid (EDTA) at 5mM, an aspartic protease inhibitor (pepstatin A at 10 μM and 20 μM and 2-mercaptoethanol at 5mM, were tested.

II. 6. Biochemical properties of purified enzyme

II. 6. 1. Optimum pH and temperature

The optimum temperature determined in the 30 to 85°C range was examined at pH 6.3 of the skim milk substratum. The pH effect on the milk coagulation enzyme expressed in relative activity (%) was evaluated by adding 0.75 mL of suitable skim milk buffer solutions to make the substrate at different pH (5.5 to 8).

II. 6. 2. Effect of CaCl₂ concentration on milk clotting

The effect of CaCl₂ concentration on milk coagulation was measured by increasing concentrations of the solution of calcium chloride (0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, and 0.08 M).

II. 6. 3. Effect of enzyme concentration

The optimum raw extract concentration was determined by observing the shortest time of milk coagulation at 0.2, 0.4, 0.6, 0.8, and 1 mg/mL.

III. Results and Discussion


III. 1. Identification of the strain

For further characterization, the PCR product of strain FK6A was examined by electrophoresis, purified and then sequenced by Geno- screen (Campus de l'Institut Pasteur de Lille, France). In addition to biochemical and morphological identification, the 16S rDNA gene was successfully amplified from the genomic DNA of the strain FK6A (**figure 1a, b**). The nucleotide database has shown 100% homology with *B. velezensis* strain. Therefore, our strain was designated *B. velezensis* FK6A with accession number MH368142 and the data bases allowed us to elaborate a phylogenetic dendrogram for FK6A and related strains (**figure 2**) according to the Neighbor Joining method by MEGA 6 software.

```

CGTCCTATCTCTAGGATTGTCAGAGGATGTCAAGACCTGGTAAGGTTCTTCGCGTTCGCTCGAATTA
ACCACATGCTCCACCGCTTGTGCGGGCCCCCGTCAATTCCTTTGAGTTTCAGTCTTGCACCGTACTC
CCCAGGCGGAGTGCTTAATGCGTTAGCTGCAGCACTAAGGGGCGGAAACCCCTAACACTTAGCACT
CATCGTTTACGGCGTGGACTACCAGGGTATCTAATCCTGTTTCGCTCCCCACGCTTTCGCTCCTCAGC
GTCAGTTACAGACCAGAGAGTGCCTTCGCCACTGGTGTTCCTCCACATCTCTACGCATTTACCGGT
ACACGTGGAATTCACCTCTCCTTCTGCACTCAAGTTCCCGAGTTTCCAATGACCCCTCCCGGTTGA
GCCGGGGCTTTCACATCAGACTTAAGAAACCGCTGCGAGCCCTTACGCCAATAATTCCGGACA
ACGCTTGCCACCTACGTATTACCGCGGTGCTGGCACGTAGTTAGCCGTGGCTTCTGTTAGGTAC
CGTCAAGGTGCCGCCCTATTTGAACGGCACTTGTCTTCCCTAACAAACAGAGCTTTACGATCCGAAAA
CCTTCATCACTACGCGGCGTTCCTCCGTGACTTTCGTCCATTGCGGAAGATCCCTACTGCTGCC
TCCCGTAGGAGTCTGGGCCGTGCTCAGTCCAGTGTGGCCGATCACC
    
```

Figure 1 (a). 16S rDNA sequence of strain FK6A with query length of 724 bp



(b). Phylogenetic tree derived from analysis of 16S rDNA sequence of strain FK6A

III. 2. Production of raw enzymatic extract

Based on the preliminary study of fermentation conditions, the production profile of enzymatic activity by the strain of *Bacillus velezensis* FK6A was found to be of interest to submerged fermentation and to grow and secrete coagulating enzymes in a short period of time (36 h). After fermentation accompanied by cold centrifugation, a raw enzyme extract was recovered by 76.6 mg/mL of protein content. A relatively lower protein concentration (11.14 mg/mL) was obtained from *Bacillus velezensis* FK6A compared to that found from *Bacillus mojavensis* P47 M in a previous study [15].

It should be mentioned also that the content reflects the total protein. Separation of the various proteins during purification will measure the concentration of required coagulating proteins. The raw extract

coagulating force is in the range of 122.05 to 3.29 US and appears to be lower compared to *Bacillus amyloliquefaciens* D4 (4645.16 US [14]) *Bacillus subtilis* B1 (782US) [7], and *Bacillus sp.*P45 (239US) [16].

The proteolytic activity obtained (0.362 ± 0.001) is relatively high, but similar to that obtained from *Bacillus subtilis* varnatto (0.440 ± 0.009) and *Rhizopus oligosporus* (0.392 ± 0.001) by Shieh et al [10]. The AC/AP obtained by the REE is also in the order of 338.12 and indicates a high proteolytic activity.

III. 3. Raw enzymatic extract purification and molecular weight estimation

Ammonium sulphate is used in the raw extract for hydrolysis and precipitation of proteins. The

gradient of precipitation used is within the precipitation of saturated enzyme proteins (80%) (**Table 1**) is consistent with that reported by Khan et al [17], Esawy and combet blanc[18] and Won Park et al [19], with enzymatic extracts of *Mucorpusillus*, *Bacillus licheniformis* 5A1 and *Bacillus sp* k-295G-7 respectively.

On Sephadex filtration gel G75, the chromatographic profile of the raw coagulating extract shows three (3) distinct absorption peaks (**Figure 2**). Clotting tests have confirmed that the first peak is inactive, while the 2nd and 3rd peaks have an enzymatic activity from the 10th to the 18th fraction, with 17.6 mL as total elution volume. The other fractions, as reported in *Bacillus amyloliquefaciens* by Ding et al [20] and in *Bacillus sphaericus*[21], represent non-active proteins

concentration range of 40 to 80%. Total probably derived from the fermentation environment. The purification parameter results were presented in **Table 2**. The fraction coagulating force is in the range of 16.45US, coagulating frequency is 0.068UP, and protein content is 76.6 mg/ml. Gel filtration chromatography is well developed as an effective technique for estimating the molecular mass of proteins.

Tableau 1. Precipitation of the crude enzyme extract with ammonium sulphate

Saturation rate (NH ₄) ₂ SO ₄ (%)	Enzyme activity in supernatant (SU)	Enzyme activity in pellet (SU)
40	17,51	10,2
60	15,13	18,7
80	-	21,76




Figure 2. Size-exclusion chromatography through Sephadex G-75 of a crude extract of *B. velezensis* FK6A strain after precipitation with 40- 80% saturated ammonium sulphate (Column 1,5 × 30 cm, Acetate buffer 0,01 M, pH 5,5, Flow rate: 36 ml/h, fraction of 2,2 ml) (■ Absorbance 280 nm, ▲ Relative activity (%))

Table 2. Purification steps of milk clotting enzyme from *B. velezensis* FK6A

Purification steps	Protein (mg/ml)	Total activity (SU)	Specific activity (U/mg protein)	Purification fold	Recovery (%)
Crude extract	76.6	122.05	1.59	1	100
(NH ₄) ₂ SO ₄ (40% - 80%).	2.34	30.76	13.14	8.26	25.20
Size exclusion Sephadex G-75	0.268	16.45	61.38	38.60	13.48

Our findings (**Figure 3**) indicate about 46 kDa of molecular weight assessed by molecular exclusion. This mass appears to be greater from *Bacillus licheniformis* strain USC13 (35 kDa) reported by Ageitos et al [6] With *Bacillus subtilis* MTCC 10422 and using the

same method, Kumari et al. [22] observed a mass of 27 kDa. Previously Yasuda et al [23] published similar findings by filtration gel followed by electrophoresis, showing a 29 kDa monomeric band. Our results are in a range of 27 to 44 kDa and consistent with the PM of several proteases from *Bacillus* sp. as reported by Peng et al [24]




Figure 3: Size-exclusion chromatography through Sephadex G-75 of a purified extract of *B. velezensis* FK6A strain (active fraction) and molecular weight determination of the milk-clotting enzyme (in box). Molecular-weight standards: 1: Bovin serum albumin (67 kDa), 2: Ovalbumin (40kDa), 3: Carbonic nhydrase (30kDa), 4: trypsin (23.8k Da) (a). **III. 4.**

Effect of some inhibitors on coagulating activity

Two protease inhibitors were tested in order to continue the biochemical characterization of the raw *Bacillus velezensis* FK6A extract and to classify the category of coagulating proteases. **Table 3** shows that pepstain A inhibits less than 13.7% of the enzyme at a concentration of 0.1 mmol/L and 48.36% at a concentration of 0.2 mmol/L. Thus, pepstain A is not to be considered

as a possible inhibitor. However, with EDTA at 5 and 10 mmol/L, coagulating activity is completely inhibited (100%), suggesting that the enzyme studied belongs to the metalloprotease class with the divalent metal ions at their active site. This group, includes mainly microbial proteases from *Bacillus amyloliquefaciens* D4 [14], *Bacillus amyloliquefaciens* JNU002 [20], *Bacillus subtilis* YB-3 ([25] and *Paenibacillus* spp. BD3526 [26]

Tableau 3. Effect of inhibitors on milk clotting activity of purified extract from *Bacillus velezensis* FK6A

Inhibitors	Concentration (mM)	Residual activity (%)
Control	0	100
Pepstain A	0,1	86,30
	0,2	51,64
2-mercaptoethanol	5	36
EDTA	5	0

III. 5. Optimal conditions for coagulating activity

III. 5. 1. Influence of temperature and pH on raw enzymatic extract (REE) activity

The optimal milk temperature on the coagulating activity of the REE of *Bacillus velezensis* FK6A strain was tested at temperatures ranging from 30 to 85°C by a level of 5°C. The findings recorded in **Figure 4A** indicate that enzymatic activity rises (2.40 ± 0.36 UP) with maximum at around 75°C. There is a significant decrease in coagulating activity after 75°C, and complete enzyme inhibition at 85°C. The optimum temperature obtained by the enzyme extract is similar to that reported by *bacilluslicheniformis* 5A5 strain [27] and close to that reported in *Bacillus subtilis* B1 (70°C) [7].

The effect of milk pH on the coagulating activity of the raw enzyme extract was measured at a pH interval of 5.7 to 8.1, with a pallier of 0.2. As shown in **Figure 4B**, 5.7 is the optimal pH of casein hydrolysis with a coagulating levels of 0.7 ± 0.01 UP. Beyond this pH value, the coagulating activity slowly decreases until total inhibition at pH 8.1. This agrees with the optimum coagulating function of the enzyme obtained from

the strain *Bacillus amyloliquefaciens* D4 with 5.5 as optimal pH and 8 as pH inactivating pH. In rennet activity, Mahaut et al [28] indicated an optimum at pH 5.8 and inactivation at pH 7.5 and complete denaturation at pH 8.

III. 5. 2. Effect of CaCl₂ concentration

Glantz et al [29] found that the calcium ion concentration in milk is linked to its gelling properties. Higher concentration in this element contributes to shorter gelling times and improved resistance and firmness of gels. The results shown in the **Figure 4C** indicate that the protease coagulating activity decreases with concentration of CaCl₂ until an optimum value of 40 mM. The same finding was reported by Talantikite et al [8] in a related analysis on *Bacillus mojavensis* I33M coagulase.

III. 5. 3. Effect of raw enzyme extract concentration

The milk clotting activity of rennet gradually increases with its concentration [30, 31]. This is in accordance with our findings (**Figure 4D**) with a steady increase in activity as the extract concentrations increase.




Figure 4. Physico-chemical properties of milk-clotting activities of the enzymatic extract from *Bacillus velezensis* FK6A (A: Temperature, B: pH, C: CaCl₂, D: Enzyme concentration)

IV. Conclusion

This strain known to generate proteolytic enzymes was not the target on its ability to coagulate milk of any research in the reviewed literature. We took an interest in this aspect by applying more rigorous methods, and by showing the originality of the concept of our

subject through the results obtained. Therefore, the proteolytic activity of *Bacillus velezensis* FK6A extracts found on the curds occurs a few minutes after the formation of a firm coagulum resulting in the presence of an abundant serum, during the technique

studies. Nevertheless, in order to solve the problem caused by The REE high proteolytic activity and totally inhibit the general proteolysis, we directed the

development towards a form of hard cooked (Figure5).



Figure 5. Appearance of cooked cheese dough making by commercial rennet (A), (B) crude extract of *Bacillus velezensis* FK6A (B) and *Bacillus velezensis* + rennet (50/50) (C)

All the results of our study show that the protease of significant coagulating potential and provides new complementary research avenues for development and

milk coagulating microbial enzymes studied has production of milk coagulating microbial enzymes.

Acknowledgements

The lab facilities provided by the department of food technology laboratory (Boumerdes University, Algeria) and the department of food technology and

human nutrition (ENSA, Algiers – Algeria) are fully acknowledged.

V. References

- Jacob M.;Jaros D.;Rohm H.; Recent advances in milk clotting enzymes. *International Journal of Dairy Technology* 64(1) (2011) 14-33.
- Mazorra-Manzano M.A.;Perea-Gutierrez T.C.;Lugo-Sanchez M.E.;Ramirez-Suarez J.C.;Torres-Llanez M.J.;Gonzalez-Cordova A.F.; Comparison of the milk-clotting properties of three plant extracts. *Food Chemistry* 141(3) (2013) 1902-7.
- Dutt K.;Meghwanshi G.K.;Gupta P.;Saxena R.K.; Role of casein on induction and enhancement of production of a bacterial milk clotting protease from an indigenously isolated *Bacillus subtilis*. *Letter of Applied Microbiology* 46(5) (2008) 513-8.
- Singh T.K.;Drake M.A.;Cadwallader K.R.; Flavor of Cheddar Cheese: A Chemical and Sensory Perspective. *Comprehensive reviews in food science and food safety* 2(4) (2003) 166-89.
- Wesley, M.Cg.; Cheddar Cheese Flavor. A Review of Current Progress. *Journal of Agriculture and Food Chemistry*, 23(6) (1975) 1047-50.
- Ageitos J.M.;Vallejo J.A.;Sestelo A.B.F.;Poza M.;Villa T.G.; Purification and characterization of a milk-clotting protease from *Bacillus licheniformis* strain USC13. *J Appl Microbiol* 103(6) (2007) 2205-13.
- Ding Z L.S., Gu Z, Zhang L, Zhang K, Shi G. Production of milk-clotting enzyme by *Bacillus subtilis* B1 from wheat bran. *African Journal of Biotechnology* 10(46) (2011) 9370-8.
- Talantikite-Kellil, S.; Nouani, A.; Saunier, M.; Gillmann, L.; Brahma, A.; Fazuane, F.; Isolation and identification of bacterial strain I33M producing milk-clotting enzyme: Optimization of culture parameters using response surface. *African Journal of Biotechnology* 11(100) (2012) 16594-606.
- Rao, L.K.; Mathur ,D.K.; Studies on the production of bacterial rennet in a pilot plant fermentor. *Biotechnology and bioengineering* 17(9)1975) 1349-61.
- Shieh, C.J.; Phan Thi, L.A.; Shih , I.L.;Milk-clotting enzymes produced by cult (ure of *Bacillus subtilis* natto. *Biochemical Engineering Journal* 43(1) (2009) 85-91.
- Silva ,B.L.;Geraldes , F.M.;Murari C.S.;Gomes E.;Da-Silva R. Production and characterization of a milk-clotting protease produced in submerged fermentation by the thermophilic fungus *Thermomucor indicae-seudaticae* N31. *Applied Biochemical and Biotechnology* 172(4) (2014) 1999-2011.
- Lowry , O.H.; Rosebrough, N.J.;Farr ,A.L.;Randall, R.J.; Protein measurement with the Folin phenol reagent. *J Biol Chem* 193(1) (1951) 265-75.
- Raposo ,S.; Domingos, A. ; Purification and characterization milk-clotting aspartic proteinases from *Centaurea calcitrapa* cell suspension cultures. *Process Biochemistry* 43(2) (2008) 139-44.
- He, X.;Ren, F.;Guo, H.; Zhang, W.; Song, X.;Gan B.; Purification and properties of a milk-clotting enzyme

- produced by *Bacillus amyloliquefaciens* D4. *Korean Journal of Chemical Engineering* 28(1) (2010) 203-8.
15. Kahlouche , A.; Nouani ,A.; Characterization of a milk-clotting enzyme produced by *Bacillus mojavensis* P47M strain isolated from Algerian dairy farm soil. *Research Journal of Biotechnology* 12(12) (2017) 37-45.
 16. Lemes, A.C.;Pavón ,Y.;Lazzaroni ,S.;Rozycki, S.;Brandelli, A.;Kalil, S.J.; A new milk-clotting enzyme produced by *Bacillus* sp. P45 applied in cream cheese development. *LWT - Food Science and Technology* 66 (2016) 217-24.
 17. Khan, M.R.;Blain, J.A.;Patterson, J.D.; Partial Purification of *Mucor pusillus* Intracellular Proteases. *Applied Environmental and Microbiology* 45(1) (1983) 94-6.
 18. Esawy ,M.A.;Combet-Blanc ,Y.; Immobilization of *Bacillus licheniformis* 5A1 milk-clotting enzyme and characterization of its enzyme properties. *World Journal of Microbiology and Biotechnology* 22(3) (2006) 197-200.
 19. Won Park, Y.;Kobayashi, H.;Kusakabe ,I.;Murakami, K.; Purification and Characterization of Soymilk-clotting Enzymes from *Bacillus* sp. K-295G-7. *Agricultural and Biological Chemistry* 51(9) (2016) 2343-9.
 20. Ding, Z. W.W.; Wang, B.; Ouyang , A.; Xiao, S.; Wang, Y.; Liu , S.; Ding , M.; Zhang, L.; Shi, G.;Production and characterization of milk-clotting enzyme from *Bacillus amyloliquefaciens* JNU002 by submerged fermentation. *European Food Research and Technology* 234(3) (2012) 415–21.
 21. El-Bendary, M.A.;Moharam, M.E.;Ali, T.H.; Purification and Characterization of Milk Clotting Enzyme Produced by *Bacillus sphaericus*. *Journal of Applied Sciences Research* 3(8) (2007) 695-9.
 22. Kumari Narwal, R.;Bhushan,B.;Pal, A.;Panwar, A.;Malhotra, S.; Purification, physico-chemico-kinetic characterization and thermal inactivation thermodynamics of milk clotting enzyme from *Bacillus subtilis* MTCC 10422. *LWT - Food Science and Technology* 65 (2016) 652-60.
 23. Yasuda, M.;Aoyama, M.;Sakaguchi,M.;Nakachi, K.;Kobamoto, N.; Purification and characterization of a soybean-milk-coagulating enzyme from *Bacillus pumilus* TYO-67. *Applied Microbiology and Biotechnology* 51(4) (1999) 474-9.
 24. Peng ,Y.;Yang, X.;Zhang, Y.; Microbial fibrinolytic enzymes: an overview of source, production, properties, and thrombolytic activity in vivo. *Applied Microbiology and Biotechnology* 69(2) (2005)126-32.
 25. Li, Y.;Liang, S.;Zhi, D.;Chen, P.;Su, F.;Li, H.; Purification and characterization of *Bacillus subtilis* milk-clotting enzyme from Tibet Plateau and its potential use in yak dairy industry. *European Food Research and Technology* 234(4) (2012) 733-41.
 26. Hang F.;Wang Q.;Hong Q.;Liu P.;Wu Z.;Liu, Z.; Zhang, H.; Chen, W.; Purification and characterization of a novel milk-clotting metalloproteinase from *Paenibacillus* spp. BD3526. *International Journal of Biological Macromolecules* 85 (2016) 547-54.
 27. Ahmed S.A.;Helmy W.A. Comparative evaluation of *Bacillus licheniformis* 5A5 and *Aloe variegata* milk-clotting enzymes. *Brazilian Journal of Chemical Engineering* 29(1)201269 - 76. v
 28. Mahaut, M.;Jeantet, R.;Brule, G;. Initiation à La Technologie Fromagère. *LAVOISIER / TEC ET DOC*. 2007.
 29. Glantz, M.;Lindmark, Mansson, H.;Stalhammar, H.;Paulsson ,M.; Effect of polymorphisms in the leptin, leptin receptor, and acyl-coenzyme A:diacylglycerol acyltransferase 1 (DGAT1) genes and genetic polymorphism of milk proteins on cheese characteristics. *Journal of Dairy Science* 4(7) (2011) 3295-304.
 30. Garnot, P.;Martin, P.; La présure: composition, activité, son rôle en fromagerie. *Technique laitière* 930(3) (1980) 27-30.
 31. Sbodio, O.A.;Tercero, E.J.;Coutaz, V.R.;Luna, J.A.;Martínez, E.; Simultaneous interaction of pH, CaCl₂ addition, temperature and enzyme concentration on milk coagulation properties / Interacción simultánea del pH, CaCl₂ añadido, temperatura y concentración enzimática sobre la coagulación de la leche. *Food science and technology international* 3(4) (2016) 291-8.

Please cite this article as:

Boumediene F., Talantikite S., Haddad S., Nouani A., Triki S., Belbraouet S., Valorisation of microbial biodiversity: Characterization of a milk-clotting enzyme produced by *Bacillus velezensis* FK6A strain isolated from Algerian dairy farm soil in light of its use in the manufacture of hard cooked cheeses. *Algerian J. Env. Sc. Technology*, 7:2 (2021) 1964-1972